


PECARN Pediatric Minor Head Injury CT Guidelines for Children Age 2+ Years



**Post-traumatic seizure IS considered to be altered mental status (AMS) and is high risk for ci-TBI based on planned secondary analyses of PECARN data*

Badawy, M. K., et al. Prevalence of Brain Injuries and Recurrence of Seizures in Children with Posttraumatic Seizures. *Academic Emergency Medicine*. 2017;24(5):595-605.

Ballard DW, Kuppermann N, Vinson DR, et al. Implementation of a Clinical Decision Support System for Children With Minor Blunt Head Trauma Who Are at Nonnegligible Risk for Traumatic Brain Injuries. *Annals of Emergency Medicine*. 2019;73(5):440-451.

Dayan PS, Holmes JF, Atabaki S, et al. Association of Traumatic Brain Injuries With Vomiting in Children With Blunt Head Trauma. *Annals of Emergency Medicine*. 2014;63(6):657-665.

Dayan PS, Holmes JF, Hoyle J, et al. Headache in Traumatic Brain Injuries From Blunt Head Trauma. *Pediatrics*. 2015;135(3):504-512.

Dayan PS, Holmes JF, Schutzman S, et al. Risk of Traumatic Brain Injuries in Children Younger than 24 Months With Isolated Scalp Hematomas. *Annals of Emergency Medicine*. 2014;64(2):153-162.

Hess EP, Homme JL, Kharbanda AB, et al. Effect of the Head Computed Tomography Choice Decision Aid in Parents of Children With Minor Head Trauma. *JAMA Network Open*. 2018;1(5).


Kuppermann N, Holmes JF, Dayan PS, et al. Identification of children at very low risk of clinically-important brain injuries after head trauma: a prospective cohort study. *The Lancet*. 2009;374(9696):1160-1170.

Lee LK, Monroe D, Bachman MC, et al. Isolated Loss of Consciousness in Children With Minor Blunt Head Trauma. *JAMA Pediatrics*. 2014;168(9):837. doi:10.1001/jamapediatrics.2014.361.

Nigrovic LE, Lee LK, Hoyle J, et al. Prevalence of Clinically Important Traumatic Brain Injuries in Children With Minor Blunt Head Trauma and Isolated Severe Injury Mechanisms. *Arch Pediatr Adolesc Med*. 2012;166(4):356-361.

Varano P, Cabrera KI, Kuppermann N, Dayan PS. Acute outcomes of isolated cerebral contusions in children with Glasgow Coma Scale scores of 14 to 15 after blunt head trauma. *Journal of Trauma and Acute Care Surgery*. 2015;78(5):1039-1043.

PECARN Pediatric Minor Head Injury CT Guidelines for Children Age <2 Years



*Post-traumatic seizure IS considered to be altered mental status (AMS) and is high risk for ci-TBI based on planned secondary analyses of PECARN data

Badawy, M. K., et al. Prevalence of Brain Injuries and Recurrence of Seizures in Children with Posttraumatic Seizures. *Academic Emergency Medicine*. 2017;24(5):595-605.

Ballard DW, Kuppermann N, Vinson DR, et al. Implementation of a Clinical Decision Support System for Children With Minor Blunt Head Trauma Who Are at Nonnegligible Risk for Traumatic Brain Injuries. *Annals of Emergency Medicine*. 2019;73(5):440-451.

Dayan PS, Holmes JF, Atabaki S, et al. Association of Traumatic Brain Injuries With Vomiting in Children With Blunt Head Trauma. *Annals of Emergency Medicine*. 2014;63(6):657-665.

Dayan PS, Holmes JF, Hoyle J, et al. Headache in Traumatic Brain Injuries From Blunt Head Trauma. *Pediatrics*. 2015;135(3):504-512.

Dayan PS, Holmes JF, Schutzman S, et al. Risk of Traumatic Brain Injuries in Children Younger than 24 Months With Isolated Scalp Hematomas. *Annals of Emergency Medicine*. 2014;64(2):153-162.

Hess EP, Homme JL, Kharbanda AB, et al. Effect of the Head Computed Tomography Choice Decision Aid in Parents of Children With Minor Head Trauma. *JAMA Network Open*. 2018;1(5).

Kuppermann N, Holmes JF, Dayan PS, et al. Identification of children at very low risk of clinically-important brain injuries after head trauma: a prospective cohort study. *The Lancet*. 2009;374(9696):1160-1170.

Lee LK, Monroe D, Bachman MC, et al. Isolated Loss of Consciousness in Children With Minor Blunt Head Trauma. *JAMA Pediatrics*. 2014;168(9):837. doi:10.1001/jamapediatrics.2014.361.

Nigrovic LE, Lee LK, Hoyle J, et al. Prevalence of Clinically Important Traumatic Brain Injuries in Children With Minor Blunt Head Trauma and Isolated Severe Injury Mechanisms. *Arch Pediatr Adolesc Med*. 2012;166(4):356-361.

Varano P, Cabrera KI, Kuppermann N, Dayan PS. Acute outcomes of isolated cerebral contusions in children with Glasgow Coma Scale scores of 14 to 15 after blunt head trauma. *Journal of Trauma and Acute Care Surgery*. 2015;78(5):1039-1043.